

Medicinal Plant Images

Ian Edwin Cock^{1,2,*}

¹Centre for Planetary health and Food Security, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland, AUSTRALIA.

²School of Environment and Science, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland, AUSTRALIA.

Correspondence:

Dr. Ian Edwin Cock

¹Centre for Planetary Health and Food Security, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.

²School of Environment and Science, Nathan Campus, Griffith University, 170 Kessels Rd, Nathan, Brisbane, Queensland 4111, AUSTRALIA.

Email id: i.cock@griffith.edu.au




Figure 1: Australian *Acacia* spp. The genus *Acacia* (family Fabaceae) is a large genus of more than 1200 trees and shrubs which are widely distributed throughout the world, with more than 700 species indigenous to Australia. The Australian species had multiple medicinal uses by indigenous Australians, including being used to treat diarrhoea and hyperglycemia^{1,2} and as a general antiseptic agent.³⁻⁷ Many Australian *Acacia* species have been reported to have antimicrobial, molluscicidal, antihypertensive and platelet aggregatory activities.¹ This photograph was taken in Brisbane, Australia in 2025 by Dr Ian Cock.



Figure 2: *Syzygium leuhmannii* (F.Muell.) L.A.S. Johnson (commonly known as riberry). *Syzygium* is a large genus of evergreen flowering plants of the family Myrtaceae which consists of approximately 500 species.⁸ Plants of this genus are widespread, occurring in tropical and subtropical regions of South-East Asia, Australia and Africa. Many *Syzygium* species produce edible fruits and berries. Some *Syzygium* spp. are used in traditional medicine to treat respiratory ailments, tuberculosis, gastro-intestinal disorders, diarrhoea and dysentery.¹ Recent reports have also highlighted the Australian *Syzygium* species *S. leuhmannii* and *Syzygium australe* (Bush Cherry) extracts as having exceptionally high antioxidant contents.⁹ Antioxidants have been associated with the prevention of cancer, cardiovascular disease and neurological degenerative disorders.¹⁰⁻¹² They are also linked with anti-diabetic bioactivities and have been associated with the reduction of obesity. Antioxidants can directly scavenge free radicals, protecting cells against oxidative stress related damage to proteins, lipids and nucleic acids.¹³ Thus, *Syzygium* spp. have potential in the treatment of a significant number of diseases and medical conditions related to cellular redox state.

Many other *Syzygium* species internationally also have documented uses in traditional medicine.² In the commercially most important species *Syzygium aromaticum* (clove), the unopened flower bud is used as a spice. This plant also has uses in traditional medicine due to its anaesthetic properties.¹⁴ The antibacterial activity of *S. aromaticum* is also well known. Numerous studies have reported on the antibacterial¹⁴ and antifungal¹⁵ activities of oils and extracts from this plant. Other *Syzygium* species from South East Asia (*Syzygium jambos*)¹⁶ and India (*Syzygium lineare* and *Syzygium cumini*)¹⁷ have also been shown to have antimicrobial activity. Recent studies have also reported the antibacterial activity of *Syzygium cordatum* leaf extracts. Of particular interest was the potent growth inhibitory of the extracts against the bacterial triggers of rheumatoid arthritis¹⁸ and ankylosing spondylitis.¹⁹

REFERENCES

1. Cock IE. Medicinal and aromatic plants – Australia. In: Ethnopharmacology, Encyclopedia of Life Support Systems (EOLSS), Developed under the auspices of UNESCO. Oxford, UK: EOLSS Publishers. Available from: <http://www.eolss.net>.
2. Cock IE, Cheesman MJ. *Acacia* spp. for the control of blood sugar in type-2 diabetes mellitus. In: Ancient and traditional foods, plants, herbs and spices in human health. Abingdon: Routledge Publishers; 2023.
3. Guo R, Yang X, Cock IE. Antimicrobial activity of *Acacia dispartima* Benth. and *Acacia leiocalyx* Pedley Leaf Extracts in Combination with Antibiotics against Bacterial Triggers of Selected Autoimmune Diseases. *Pharmacogn Commun.* 2023;13(3):104-15. doi: 10.5530/pc.2023.3.18.
4. Cock IE. Antibacterial activity of selected Australian native plant extracts. *Internet J Microbiol.* 2008;4:2.
5. Cock IE. Antimicrobial activity of *Acacia aulacocarpa* and *Acacia complanta* methanolic extracts. *Pharmacogn Commun.* 2012;2(1):66-71. doi: 10.5530/pc.2012.1.12.
6. Cock IE. Australian *Acacia* spp. extracts as natural food preservatives: growth inhibition of food spoilage and food poisoning bacteria. *Pharmacogn Commun.* 2017;7(1):4-15. doi: 10.5530/pc.2017.1.2.
7. Cock IE, Winnett V, Sirdaarta J, Matthews B. The potential of selected Australian medicinal plants with anti-Proteus activity for the treatment and prevention of rheumatoid arthritis. *Pharmacogn Mag.* 2015;11(42) Suppl 1:S190-208. doi: 10.4103/0973-1296.157734, PMID 26109767.
8. Cock IE, Cheesman M. Plants of the genus *Syzygium* (Myrtaceae): a review on ethnobotany, medicinal properties, and phytochemistry. In: Bioactive compounds of medical plants. Properties and potential for human health. NJ: Apple Academic Press; 2018.
9. Netzel M, Netzel G, Tian Q, Schwartz S, Konczak I. Native Australian fruits – a novel source of antioxidants for food. *Innov Food Sci Emerg Technol.* 2007;8(3):339-46. doi: 10.1016/j.ifset.2007.03.007.
10. Potter JD. Cancer prevention: epidemiology and experiment. *Cancer Lett.* 1997;114(1-2):7-9. doi: 10.1016/S0304-3835(97)04615-6, PMID 9103244.
11. Cock IE. The medicinal properties and phytochemistry of plants of the genus *Terminalia* (Combretaceae). *Inflammopharmacology.* 2015;23(5):203-29. doi: 10.1007/s10787-015-0246-z, PMID 26226895.
12. Cock IE. Problems of reproducibility and efficacy of bioassays using crude extracts, with reference to *Aloe vera*. *Pharmacogn Commun.* 2011;1(1):52-62. doi: 10.5530/pc.2011.1.3.
13. Rice-Evans C, Miller N, Paganga G. Antioxidant properties of phenolic compounds. *Trends Plant Sci.* 1997;2(4):152-9. doi: 10.1016/S1360-1385(97)01018-2.
14. Arora DS, Kaur GJ. Antibacterial activity of some Indian medicinal plants. *J Nat Med.* 2007;61(3):313-7. doi: 10.1007/s11418-007-0137-8.
15. Park MJ, Gwak KS, Yang I, Choi WS, Jo HJ, Chang JW *et al.* Antifungal activities of the essential oils of *Syzygium aromaticum* (L.) Mer. Et Perry and *Leptospermum petersonii* Bailey and their constituents against various dermatophytes. *J Microbiol.* 2007;45(5):460-5. PMID 17978807.
16. Mohanty S, Cock IE. Bioactivity of *Syzygium jambos* methanolic extracts: antibacterial activity and toxicity. *Pharmacogn Res.* 2010;2(1):4-9. doi: 10.4103/0974-8490.60577, PMID 21808530.
17. Duraipandiyan V, Ayyanar M, Ignacimuthu S. Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. *BMC Complimentary Altern Med.* 2006;6(35):1-7.
18. Cock IE, van Vuuren SF. Anti-Proteus activity of some South African medicinal plants: their potential for the prevention of rheumatoid arthritis. *Inflammopharmacology.* 2014;22(1):23-36. doi: 10.1007/s10787-013-0179-3, PMID 23877712.
19. Cock IE, Van Vuuren SF. The potential of selected South African plants with anti-Klebsiella activity for the treatment and prevention of ankylosing spondylitis. *Inflammopharmacology.* 2015;23(1):21-35. doi: 10.1007/s10787-014-0222-z, PMID 25412961.