

The relaxant effect induced by *Allium sativum* L. bulb aqueous extract on rat isolated trachea

Badreddine Fehri, Mueen Ahmed K.K.¹, Jean-Marc Aiache²

Department of Pharmacology and Toxicology, Tunisia Pharmaceutical Industries Company (SIPHAT), Fondouk Choucha-Radès, 2013 Ben Arous, Tunisia, ¹Department of Pharmaceutical Sciences, College of Clinical Pharmacy, King Faisal University, P. O. Box 400, Al-Ahsa-31982, Kingdom of Saudi Arabia, ²Department of Biopharmaceutics, Faculty of Pharmacy, Auvergne University, 28, Place Henri Dunant - BP 38, 63001 Clermont-Ferrand, France

Submitted: 16-10-2010

Revised: 10-11-2010

Published: 20-01-2011

ABSTRACT

Background: Garlic plays an important role in complementary and alternative medicine. Most people believe in and use herbal products even when they have not been as thoroughly researched as garlic. Garlic is also known for its beneficial effects on the cardiovascular system. **Materials and Methods:** The relaxant effect of *Allium sativum* L. bulb aqueous extract (ASBAE) containing 0.06%-0.10% of allicin was studied on isolated smooth muscle of trachea of rats precontracted using acetylcholine (10^{-5} M). **Results:** It was found that ASBAE induced a dose-dependent relaxation with recorded EC_{50} values of $71.87 \pm 5.90 \mu\text{g/mL}$ ($n = 7$). Pretreatments with mepyramine (10^{-7} M), methysergide (10^{-7} M), caffeine (10^{-6} M), theophylline (10^{-6} M), nifedipine (10^{-6} M), and dipyridamole (10^{-6} M) did not alter ASBAE concentration-response curves. In turn, concentration-response curves to ASBAE were significantly shifted toward right in the presence of aspirin (3.10^{-3} M), indomethacin (10^{-6} M), prazosin (10^{-6} M), and propranolol (10^{-7} M). **Conclusion:** It is suggested that the recorded relaxation results are due to the release of prostaglandins E_1 and E_2 consecutively to α - and β -adrenoreceptor stimulation.

Key words: *Allium sativum* L. bulb aqueous extract, rat isolated trachea

Access this article online

Website:

www.phcog.com

DOI:

10.4103/0973-1296.75879

Quick Response Code:



INTRODUCTION

Allium sativum, commonly known as garlic, is one of the species belonging the onion family Alliaceae. Garlic is largely used around the world for its pungent flavor as a seasoning or condiment.^[1] It is a fundamental component in many or most dishes of various regions, including eastern Asia, south Asia, Southeast Asia, the Middle East, northern Africa, southern Europe, and parts of South and Central America.^[2] It contains the sulfur-containing compounds alliin, ajoene, diallylsulfide, dithiin, S-allyl cysteine and also enzymes, vitamin B, proteins, minerals, saponins, flavonoids, and maillard reaction products, which are non-sulfur-containing compounds. Furthermore, a phytoalexin called allixin (3-hydroxy-5-methoxy-6-methyl-2-pentyl-4H-pyran-4-one) was found, a nonsulfur compound with

a γ -pyrone skeleton structure with antioxidative effects, antimicrobial effects, antitumor promoting effects, inhibition of aflatoxin B₂ DNA binding, and neurotrophic effects.^[3-8]

A. sativum L. was also demonstrated to reduce cardiovascular risk factors, especially blood pressure,^[9-11] hyperlipidemia, and atherosclerosis.^[12] Garlic is also shown to have significant relaxant effects on the airway smooth muscles. In this regard, it has been observed that administering garlic prevented the occurrence of high altitude breathing symptoms.^[13] In 1998, Fallon *et al.*^[14] demonstrated that active substances that are found in garlic caused vasorelaxation in the pulmonary arterial segments. Recently, a report indicated that a mixture of herbs containing *A. sativum* L. reduced an elevated respiratory rate in horses with recurrent airway obstruction.^[15] In contrast, the mechanism of action of *A. sativum* L. in airway smooth muscle remains unclear. Hence, the present study is to investigate and identify the possible mechanism for the relaxant effects caused by *A. sativum* L. extracts.

Address for correspondence:

Dr. Badreddine Fehri, B.P. 181, 2040 Radès, Tunisia.
E-mail: bms.fehri@yahoo.com

MATERIALS AND METHODS

Plant material

The drug studied was a *A. sativum* L. bulb aqueous extract (ASBAE) obtained from Finzelberg Laboratories (Germany), which answered the following parameters:

- Ratio of dried plant/extract: 6:1
- Allicin content was determined using gas chromatography: 0.10%

Drugs used

ASBAE (Finzelberg), acetylcholine HCl, propranolol HCl, dipyridamole were purchased from Sigma, St. Louis, MO, USA, nifedipine from Dolder, Basel, Switzerland, aspirin DL-lysine (DL-lysine acetylsalicylate) from Sanofi-aventis Paris, France, prazosin HCl from Pfizer Kalamazoo, USA, caffeine, theophylline sodium anisate, methysergide maleate, indomethacin, and mepyramine maleate from Siphat, Fondouk Choucha-Radès, Tunisia.

Prior to their dilution in Tyrode's solution, indomethacin, dipyridamole, and nifedipine were dissolved in absolute ethanol and then diluted in Tyrode's solution. Theophylline and aspirin were used from ampoules for injection (Siphat, Fondouk Choucha-Radès, Tunisia and Sanofi-aventis, Paris, France). The other substances were directly dissolved in Tyrode's solution. The required concentrations of the compounds were added to the organ bath so that the volume of ethanol never exceeded 0.05 mL (0.4% by volume). Such quantity of ethanol had no effect on acetylcholine and ASBAE responses on isolated rat trachea.

Rat trachea in vitro

Male rats weighing 300-400 g were selected in a homogeneous breeding center (Central Animal House, Tunisia Pharmaceutical Industries Company, Fondouk Choucha-Radès, Tunisia). They were sacrificed by a blow and exsanguinated. The trachea was removed immediately and cut into small segments of 3-4 rings, which were equilibrated under a resting tension of 1.5 g and suspended in 10 mL baths containing Tyrode's solution at $37^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$ and the tissue was aerated continuously with 95% O_2 and 5% CO_2 . The composition of the Tyrode's solution was (mM): [NaCl: 139.2; KCl: 2.7; CaCl_2 : 1.8; MgCl_2 : 0.49; NaHCO_3 : 11.3; Na_2HPO_4 : 0.4; glucose: 5.5]. Tension was measured isometrically with Ugo Basil strain gauges displayed on Gemini Ugo Basile recorders (Italy).

Experimental procedure

The tracheal rings (3 to 4) were placed in organ bath for 85 min and washed with Tyrode solution at every 15-min time interval. The tissue was contracted using acetylcholine (ACh 3.10^{-3} M) until maximal contraction was achieved. During that stage, the tissue was subjected to relaxation

using with theophylline (THEO 3.10^{-3} M), further washed repeatedly and allowed to rest until it returned to baseline tension. When baseline tension was reached, it was again contracted with ACh 10^{-5} M, and maximal tension was obtained, during this time, it was studied to establish concentration-response curves to ASBAE by the method of cumulative addition. The final addition was followed by addition of THEO 3.10^{-3} M to obtain complete relaxation. Pretreatments with drugs were performed 30 min before ACh 10^{-5} M contractions. The relaxations induced by ASBAE were expressed as a percentage of relaxation vs complete relaxation (E_{MAX}). The $-\log \text{EC}_{50}$ (pD_2) values, defined as the negative log concentration-effect curves, were evaluated graphically from each experiment.

Statistical analysis

All data were expressed as mean $\pm$ SEM. Data of relaxant effects of different concentrations of extracts were compared with the results of negative and positive control using Student *t* test. Significant differences with controls are shown as $*P < 0.05$ and $***P < 0.001$.

RESULTS

ASBAE exhibited a dose-dependent relaxant effect on the rat isolated trachea precontracted with ACh 10^{-5} M [Figure 1]. On basal tone of isolated tissue, ASBAE showed mild contractile effects at lower doses ranging from 10^{-6} to 10^{-4} g/mL, followed by relaxant effects at higher doses (from 10^{-3} to 3.10^{-3} g/mL) (data not shown).

Pretreatments with mepyramine 10^{-7} M and methysergide 10^{-7} M, ASBAE concentration-response curves were not significantly modified as observed by pD_2 values




Figure 1: Representative trace from acetylcholine (ACh 10^{-5} M) precontracted rat isolated trachea showing cumulative dose-response to *Allium sativum* L. aqueous extract. Concentrations (10^{-6} to 3.10^{-3}) are in g/mL

[Table 1]. Statistical analysis of pD_2 values also failed to detect any difference in the relaxant effect of ASBAE between control and tracheal rings treated with caffeine (10^{-6} M), theophylline (3.10^{-6} M), nifedipine (10^{-6} M), and dipyridamole (10^{-6} M) [Table 1]. In turn, pretreatments with aspirin (3.10^{-3} M) and indomethacin (10^{-6} M) caused right shift to ASBAE concentration-response curves, and pD_2 values significantly decreased [Table 1 and Figure 2]. Prazosin (10^{-6} M) and propranolol (10^{-7} M) had the same effect ($P < 0.05$ and $P < 0.001$) [Table 1 and Figure 3].

DISCUSSION

The present work indicated that ASBAE induced multiple effects on rat isolated trachea according to the level on tone. On basal tone trachea, ASBAE induced contractile

Table 1: pD_2 ($-\log EC_{50}$) of ASBAE and its influence of various pretreatments on relaxant effect

Pretreatment (M)	N	pD_2
Control	7	4.14 ± 0.03
Caffeine (10^{-6})	3	4.17 ± 0.06
Theophylline (3.10^{-6})	3	4.18 ± 0.03
Mepyramine (10^{-7})	3	4.27 ± 0.15
Methysergide (10^{-7})	3	4.31 ± 0.10
Dipyridamole (10^{-6})	3	4.28 ± 0.08
Nifedipine (10^{-6})	3	4.48 ± 0.25
Aspirin (3.10^{-3})	3	$3.84 \pm 0.09^*$
Indomethacin (10^{-6})	3	$4.01 \pm 0.01^*$
Prazosin (10^{-6})	3	$4.03 \pm 0.01^*$
Propranolol (10^{-7})	3	$3.25 \pm 0.13^{***}$

ASBAE, *Allium sativum* L. bulb aqueous extract, Values are means $\pm$ SEM. The relaxations induced by ASBAE were expressed as a percentage of maximal relaxation (E_{MAX}). The pD_2 ($-\log EC_{50}$) values were derived graphically from the log concentration-effect curves; n indicates the number of tracheal preparations. Significant differences with controls are shown as: $^*P < 0.05$ and $^{***}P < 0.001$




Figure 2: Cumulative concentration-response curves for *Allium sativum* L. extract on rat isolated trachea. Values are mean $\pm$ SEM

effects at low doses (10^{-6} to 10^{-4} g/mL), whereas at higher doses (10^{-3} to 3.10^{-3} g/mL), it induced relaxation effect.

Furthermore, it was also demonstrated through the present work that ASBAE induced a dose-dependent relaxation with EC_{50} values of 71.87 ± 5.90 μ g/mL ($n = 7$) on the rat isolated trachea precontracted with ACh 10^{-5} M. In the presence of both, mepyramine 10^{-7} M, an antagonist of H_1 receptors^[16] and methysergide 10^{-7} M, an antagonist of serotonin (5-Hydroxytryptamine) receptors,^[17] the induced relaxation of ASBAE was not significantly modified. The present study suggests that ASBAE-induced relaxation may not be resulting from interactions with histamine and serotonin receptors. In the other hand, caffeine (10^{-6} M) and theophylline (3.10^{-6} M), two phosphodiesterase (PDE) inhibitors,^[18] and nifedipine (10^{-6} M) and dipyridamole (10^{-6} M), two compounds acting as calcium antagonists,^[19] did not alter ASBAE-induced relaxation; hence, it is possible to suggest that the recorded relaxation does not involve inhibition of cyclic nucleotide phosphodiesterase or a calcium blocking influx phenomenon.

The prostaglandin synthetase inhibitors, aspirin (3.10^{-3} M) and indomethacin (10^{-6} M),^[19] significantly antagonized ASBAE-induced relaxation. The present outcome may possibly suggest that the recorded relaxation could be mediated due to the release of the relaxant metabolites of arachidonic acid (AA) produced under the activation of cyclooxygenase. It has been shown that garlic prevents platelet aggregation by selectively inhibiting platelet thromboxane (TX) synthesis and reducing vascular prostacyclin (PGI_2).^[20,21]

The lungs of many animal species, including man, are rich in prostaglandins (PGs) that do influence the tone of




Figure 3: Cumulative concentration curves for *Allium sativum* L. extract on rat isolated trachea. Values are mean $\pm$ SEM

tracheal and bronchial muscles. Many studies demonstrated that PGs of the E series and PGs of F series, respectively, induce a relaxation and a contraction of tracheal smooth muscle *in vitro*. Takano *et al.*^[22] particularly studied the effects of PGs E₁, E₂, F_{2α}, A₁, A₂, and B₂ on guinea pig-isolated trachea and showed that F_{2α}, B₂, and A₂ produced contraction, whereas PGs E₁ and E₂ produced relaxation. The same authors reported that PG A₁ produced no response. The same result was obtained by Coleman and Kennedy,^[23] who studied the effect of 12 PGs on the guinea pig-isolated trachea. PGs of series E are considered to be bronchodilators not only in experimental animals but also in man.^[24] Taking into consideration, based on these observations, it is now possible to suggest that active components contained in ASBAE induced the recorded relaxation by a stimulation of PGs E₁ and E₂, and further, the effect was antagonized by aspirin and indomethacin pretreatments.

It is also reported that PGs E₁ and E₂ cause relaxation when tone is present and contraction when tone is absent.^[19,24,25] This may explain the contractile effects induced at low doses (from 10⁻⁶ to 10⁻⁴ g/mL) by ASBAE on basal tone trachea.^[26] This contraction phase could therefore be mediated by PGs E₁ and E₂. When basal tone has reached a sufficient level, then a relaxation phase occurred (from 10⁻³ to 3.10⁻³ g/mL) probably due to the release of PGs E₁ and E₂.

Pretreatments with the respective α- and β-adrenoreceptor blockers, prazosin 10⁻⁶ M^[27,28] and propranolol 10⁻⁷ M, caused right shift of ASBAE concentration-effect curves. The order of potency of the recorded antagonism was propranolol 10⁻⁷ M > prazosin 10⁻⁶ M [Table 1 and Figure 3].

This result may suggest that active components present in ASBAE could stimulate both α- and β-adrenergic receptors with a higher affinity for β than for α receptors. The affinity of various *Allium* species for adrenergic (α₁, α₂, β₁, and β₂) receptors has recently been studied. Nencini *et al.*^[29] reported that *A. sativum* L. did not show high affinity for α₁ and α₂ receptors and showed a partial affinity for β₁ receptors, whereas wild-type *Allium* species presented a greater affinity for these receptors, in particular for β₂ receptors (*Allium neapolitanum* and *Allium subhirsutum* species).

CONCLUSION

Further investigations need to be undertaken to determine the affinity of *A. sativum* L. for adrenergic receptors on rat trachea. The link between PGs E₁ and E₂ and α- and β-adrenergic receptor stimulation by which active

components contained in ASBAE relax the rat isolated trachea should also be clarified. In this respect, it has been suggested that PGs E₂ exert their biologic actions via at least 4 receptor subtypes (EP₁-EP₄) and that EP₁ could act as a modulator of β₂-adrenergic receptor.^[30] Active components present in ASBAE could possibly relax the rat isolated trachea by such mechanism of action.

REFERENCES

- Gardner CD, Messina M, Lawson LD, Farkuhar JW. Soy, garlic, and *Ginkgo biloba*: Their potential role in cardiovascular disease prevention and treatment. *Curr Atheroscler Rep* 2003;5:468-75.
- Khanum F, Anilakumar KR, Viswanathan KR. Anticarcinogenic properties of garlic. A review. *Crit Rev Food Sci Nutr* 2004;44:479-88.
- Tattleman E. Health effects of garlic. *Am Fam Phys* 2005;72:103-6.
- Sharma A, Sharma V, Kansal L. Therapeutic Effects of *Allium sativum* on Lead-induced Biochemical changes in Soft tissues of Swiss Albino Mice. *Phcog Mag* 2009;5:364-71. Available from: <http://www.phcog.com/text.asp?2009/5/20/364/58568> [last cited on 2010 Oct 12].
- Navaneetha LK, Veerakumari L. Effect of *Allium sativum* on the phosphoenolpyruvate carboxykinase and pyruvate kinase activity of *Haemonchus contortus in vitro*. *Phcog Mag* 2009;5:430-2.
- Petrovska BB, Cekovska S. Extracts from the history and medical properties of garlic. *Phcog Rev* 2010;4:106-10.
- Afkhami-Ardekani M, Kamali-Ardekani A, Shojaoddiny-Ardekani A. Effects of garlic on serum lipids and blood glucose of type 2 diabetic patients. *Int J Diab Dev Ctries* 2006;26:86-8.
- Biren N Shah, Bhavesh S Nayak, Manish A Patel, Ashish D Mishra, Avinash K Seth. Aged garlic extract-a health benefit review. *Phcog Mag* 2006;2:90-5.
- Banerjee SK, Maulik SK. Effect of garlic on cardiovascular disorders: A review. *Nutr J* 2002;1:4-14.
- Dhawan V, Jain S. Effect of garlic supplementation on oxidized low density lipoproteins and lipid peroxidation in patients of essential hypertension. *Mol Cell Biochem* 2004;266:109-15.
- Turner B, Mølgaard C, Marckmann P. Effect of garlic (*Allium sativum*) powder tablets on serum lipids, blood pressure and arterial stiffness in normo-lipidaemic volunteers: A randomized, double blind, placebo-controlled trial. *Br J Nutr* 2004;92:701-6.
- WHO. *Bulbus Allii Sativi*. In: WHO monographs on selected medicinal plants. Vol 1. Geneva WHO; 1999. p. 16-32.
- Harris L. The book of Garlic. Reading, MA: Addison-Wesley; 1995.
- Fallon MB, Gary AA, Tarek TA, Jun D, Shi JC, Yiu FC, *et al.* Garlic prevents hypoxic pulmonary hypertension in rats. *Am J Physiol Lung Cell Mol Physiol* 1998;275:283-7.
- Pearson W, Charch A, Brewer D, Clarke AF. Pilot study investigating the ability of an herbal composite to alleviate clinical signs of respiratory dysfunction in horses with recurrent airway obstruction. *Can J Vet Res* 2007;78:145-51.
- Parsons ME, Ganellin CR. Histamine and its receptors. *Br J Pharmac* 2006;147:S127-35.
- Saxena PR, Lawang A. A comparison of cardiovascular and smooth muscle effects of 5-hydroxytryptamine and 8-carboxamidotryptamine, a selective agonist of HT₁ receptors. *Arch Int Pharmacodyn Ther* 1985;277:235-52.

18. Weiss B, Hait WN. Selective cyclic nucleotide phosphodiesterase inhibitors as potential therapeutic agents. *Ann Rev Pharmacol Toxicol* 1977;17:441-77.
19. Nayler WG, Poole-Wilson P. Calcium antagonists: definition and mode of action. *Basic Res Cardiol* 1981;76:1-15.
20. Vane T, Booting R. Inflammation and mechanism of action of anti-inflammatory drugs. *FASSEB J* 1987;1:89-96.
21. Ali M, Mohammed SY. Selective suppression of platelet thromboxane formation with sparing of vascular prostacyclin synthesis by aqueous extract of garlic in rabbits. *Prostaglandins Leukot Med* 1986;25:139-46.
22. Takano S, Sawano A, Watanabe H, Suzuki T. A comparison of response of guinea-pig isolated trachea to six prostaglandins. *Prostaglandins* 1978;15:485-9.
23. Coleman RA, Kennedy I. Contractile and relaxant actions of prostaglandins on guinea-pig isolated trachea. *Br J Pharmacol* 1980;68:533-9.
24. Rosenthale ME. Prostaglandins as bronchodilators. *N Y State J Med* 1975;75:374-8.
25. Puglish L. Opposite effects of prostaglandins E and F on tracheal smooth muscles and their interactions with calcium ions. *Adv Biosci* 1973;9:219-27.
26. Lambley JE, Smith AP. The effects of arachidonic acid, indomethacin and SC-19220 on guinea-pig tracheal muscle tone. *Eur J Pharmac* 1975;30:148-53.
27. Davey MJ. The pharmacology of prazosin, an alpha-adrenoreceptor antagonist and the basis for its use in the treatment of essential hypertension. *Clin Exp Hypertens* 1982;a4:1-2,47-59.
28. Frishman WH, Cheng-Lai A. Nawarskas *Current Cardiovascular Drugs*. 4th ed. *Curr Med* 2005. p. 152.
29. Nencini C, Franchi GG, Micheli L. Cardiovascular receptor binding affinity of aqueous extracts from *Allium* species. *Int J Food Sci Nutr* 2010;61:433-9.
30. McGraw DW, Mihibachler KA, Schwarb MR, Rahman FF, Small KM, Almoosa KF, *et al.* Airway smooth muscle prostaglandin-EP₁ receptors directly modulate β_2 -adrenergic receptors within a unique heterodimeric complex. *J Clin Invest* 2006;116:1400-9.

Source of Support: Nil, **Conflict of Interest:** None declared

Author Help: Reference checking facility

The manuscript system (www.journalonweb.com) allows the authors to check and verify the accuracy and style of references. The tool checks the references with PubMed as per a predefined style. Authors are encouraged to use this facility, before submitting articles to the journal.

- The style as well as bibliographic elements should be 100% accurate, to help get the references verified from the system. Even a single spelling error or addition of issue number/month of publication will lead to an error when verifying the reference.
- Example of a correct style
Sheahan P, O'leary G, Lee G, Fitzgibbon J. Cystic cervical metastases: Incidence and diagnosis using fine needle aspiration biopsy. *Otolaryngol Head Neck Surg* 2002;127:294-8.
- Only the references from journals indexed in PubMed will be checked.
- Enter each reference in new line, without a serial number.
- Add up to a maximum of 15 references at a time.
- If the reference is correct for its bibliographic elements and punctuations, it will be shown as CORRECT and a link to the correct article in PubMed will be given.
- If any of the bibliographic elements are missing, incorrect or extra (such as issue number), it will be shown as INCORRECT and link to possible articles in PubMed will be given.